

## **Environmental Mycotoxin Panel**

Company: Project Name:	Accession #: External ID:	Collected:
Address:	Phone:	Received: Completed:
E-Mail:		

**FELINE** 

**Procedure Type:** Semi-Quantitative ELISA (Enzyme-Linked Immunosorbent Assay)

Code	Test - Pet Urine	Result	Value (ppb)	Reference Range
OCHRA	Ochratoxin A	Present	2.515	<1.8 Not Present  1.8 to <2 Equivocal  >=2 Present
AFLA	Aflatoxin Group: (B1, B2, G1, G2)	Equivocal	0.938	<0.8 Not Present  0.8 to <1 Equivocal  >=1 Present
TRICHO	Trichothecene Group (Macrocyclic): Roridin A, Roridin E, Roridin H, Roridin L-2, Verrucarin A, Verrucarin J, Satratoxin G, Satratoxin H, Isosatratoxin F	Present	0.149	<0.07 Not Present 0.07 to <0.09 Equivocal >=0.09 Present
GLIO	Gliotoxin Derivative	Equivocal	0.901	<0.5 Not Present 0.5 to <1 Equivocal >=1 Present
ZEA	Zearalenone	Not Present	0.042	<0.5 Not Present 0.5 to <0.7 Equivocal >=0.7 Present

RTL maintains liability limited to cost of analysis. Interpretation of the data contained in this report is the responsibility of the client.

This report relates only to the samples reported above and may not be reproduced, except in full, without written approval by RTL.

The above test report relates only to the items tested. RTL bears no responsibility for sample collection activities or analytical method limitations.



	Mycotoxin	Cellular Activity of Mycotoxin	Symptoms/Other	Association with a "Disease State"	
			A FLATOXIN FA MILY		
	Organisms: Aspergillus flavus, Aspergillus oryzae, Aspergillus fumigatus, Aspergillus parasiticus				
		Aflatoxins have been associate	d with liver cancer [2,3], cirrhosis [4,5], and other he	alth issues	
1	Aflatoxin B1	Binds DNA and proteins [6,7]	Shortness of breath [8], weight loss [10], most potent and highly carcinogenic.	Primarily attacks the liver, other organs include kidneys and lungs [11]	
2	Aflatoxin B2	Inhibits DNA and RNA replication [12]	Impaired fetal growth [13,14]	Affects the liver and kidneys [11]	
3	Aflatoxin G1	Cytotoxic, induces apoptosis in cells, DNA damage [1]	A flavus is a leading cause of invasive aspergillus in immunocompromised patients [15]	Cancer, neonatal jaundice [2,3,16]	
4	Aflatoxin G2	Cancer, neonatal jaundice [2,3,16]	Aflatoxicosis in humans and animals [15]	Malnutrition, lung cancer [2,3,16]	
			OCHRATOXIN A		
		Organisms: <i>Asperg</i>	illus ochraceus, Aspergillus niger, Penicillium species		
5	Ochratoxin A	Inhibits mitochondrial ATP, potent teratogen, and immune suppressor [17-19]	Fatigue, dermatitis, irritated bowel [20-22]	Kidney disease and cancer [23,24]	
	MA CROCYCLIC TRICHOTHECENES (Group D)				
		C	rganism: Stachybotrys chartarum		
6	Satratoxin G	DNA, RNA, and protein synthesis inhibition [25]	Fatigue [26]	Bleeding disorders, nervous system disorders [27,28]	
7	Satratoxin H	Inhibits protein synthesis [25]	Fatigue [26]	Breathing issues [29]	
8	Isosatratoxin F	Immunosuppression [30]	Weakened immune system [30]		
9	Roridin A	Immunosuppression [30]	Weakened immune system [30]		
10	Roridin E	DNA, RNA, and protein synthesis disruption [25,32]	Weakened immune system [30]	Lung and nasal olfactory problems [31]	
11	Roridin H	Inhibits protein synthesis [25]	Weakened immune system [30]		
12	Roridin L-2	Immunosuppression [30]	Weakened immune system [30]		
13	Verrucarin A	Immunosuppression [30]			
14	Verrucarin J	Immunosuppression [30]			
	GLIOTOXIN DERIVATIVE				
	Organisms: Aspergillus fumigatus, Aspergillus terreus, Aspergillus niger, Aspergillus flavus				
15	Gliotoxin	Attacks intracellular function in immune system [34]	Memory and breathing issues [35,36]	Immune dysfunction disorders [34]	
	ZEARALENONE				
			Organisms: Füsarium species		
16	Zearalenone	Estrogen mimic [37,38]	Early puberty, low sperm counts, cancer [39-42]	Cancer [39,40]	

## REFERENCES:

- 1 Zhang, Z., et al., Cytochrome P450 2A13 is an efficient enzyme in metabolic activation of aflatoxin 61 in human bronchial epithelial cells. Arch Toxicol, 2013. 87(9): p. 1697-707.
- Wang, S.H., S.H. Yeh, and P.J. Chen, Androgen Enhances Aflatoxin-induced Genotoxicity and Inflammation to Liver Cancer in Male Hepatitis B Patients. Cell Mol Gastroenterol Hepatol, 2023. 15(2): p. 507-508.
- Fan, J.H., et al., Attributable causes of liver cancer mortality and incidence in china. Asian Pac J Cancer Prev, 2013. 14(12): p. 7251-6.
- Seitz, H.K. and F. Stickel, Risk factors and mechanisms of hepatocarcinogenesis with special emphasis on alcohol and oxidative stress. Biol Chem, 2006. 387(4): p. 349-60. Chu, Y.J., et al., Aflatoxin B(1) exposure increases the risk of circhosis and hepatocellular carcinoma in chronic hepatitis 8 virus carriers. Int Cancer, 2017. 141(4): p. 711-720.
- Lin, Y.C., et al., DNA polymerase zeta limits chromosomal damage and promotes cell survival following aflatoxin exposure. Proc Natl Acad Sci U S A, 2016. 113(48): p. 13774-13779. Poirier, M.C., Chemical-induced DNA damage and human cancer risk. Discov Med, 2012. 14(77): p. 283-8.
- Le Pape, P., et al., First case of Aspergillus caelatus airway colonization in a Chronic Obstructive Pulmonary Disease patient. Int. Infect Dis, 2019. 81: p. 85-90.

  Hernandez-Martinez, R. and I. Navarro-Blasco, Affatoxin levels and exposure assessment of Spanish Infant cereals. Food Addit Contam Part B Surveill, 2010. 3(4): p. 275-88.
- 10 Melaram, R., Environmental Risk Factor's Implicated in Liver Disease: A Mini-Review. Front Public Health, 2021. 9: p. 683719.
- 11 Pelkonen, O. and H. Raunio, Metabolic activation of toxins: tissue-specific expression and metabolism in target organs. Environ Health Perspect, 1997. 105 Suppl 4(Suppl 4): p. 767-74. 12 Madrigal-Santillan, E., et al., Antigenotoxic studies of different substances to reduce the DNA damage induced by aflatoxin B(1) and ochratoxin A. Toxins (Basel), 2010. 2(4): p. 738-57.
- 13 Tesfamariam, K., et al., Chronic aflatoxin exposure during pregnancy is associated with lower fetal growth trajectories: a prospective cohort from the Butajira Nutrition, Mental Health, and Pregnancy (BUNMAP) Study in rural Ethiopia. Am J Clin Nutr, 2022. 116(6): p. 1634-1641.

  14 Smith, L.E., et al., Aflatoxin Exposure During Pregnancy, Maternal Anemia, and Adverse Birth Outcomes. Am J Trop Med Hyg, 2017. 96(4): p. 770-776.

- 15 Sugui, J.A., et al., Assessment of serum aflatoxin B(1) levels in neonatal jaundice with glucose-6-phosphate dehydrogenase deficiency: a preliminary study. Mycotoxin Res, 2021. 37(1): p. 109-116.
- 17 Al-Anati, L. and E. Petzinger, Immunotoxic activity of ochratoxin A. J Vet Pharmacol Ther, 2006. 29(2): p. 79-90. 18 Tao, Y., et al., Ochratoxin A: Toxicity, oxidative stress and metabolism. Food Chem Toxicol, 2018. 112. p. 320-331.

- 20 Wu, T.Y., et al., Prevalence of Aspergillus-Derived Mycotoxins (Ochratoxin, Aflatoxin, and Gliotoxin) and Their Distribution in the Urinalysis of ME/CFS Patients. Int J Environ Res Public Health, 2022, 19(4).
- 21 Akiyama, T,, et al., The human cathelicidin LL-37 host defense peptide upregulates tight junction related proteins and increases human epidermal keratinocyte barrier function. J Innate Immun, 2014. 6(6): p. 739-53.
- 22 Gao, Y., et al., The Compromised Intestinal Barrier Induced by Mycotoxins. Toxins (Basel), 2020. 12(10).
- 23 Clark, H.A. and S.M. Snedeker, Ochratoxin a: its cancer risk and potential for exposure. J.Toxicol Environ Health B Crit Rev, 2006. 9(3): p. 265-96.
- 24 Fuchs, R., and M. Perajka, Octivatoxin A in human kidney diseases. Food Addit Contam, 2005. 22 Suppl 1: p. 53-7.
  25 Yang, G.H., et al., Apoptosis induction by the satratoxins and other trichothecene mycotoxins: relationship to ERK, p38 MAPK, and SAPK/JNK activation. Toxicol Appl Pharmacol, 2000. 164(2): p. 149-60. 26 Johanning, E., et al., Health and immunology study following exposure to toxigenic fungi (Stachybotrys chartarum) in a water-damaged office environment. Int Arch Occup Environ Health, 1996. 68(4): p. 207-18.
- 2° 68(4): p. 207-18.

  27 Islam, Z., et al., Purification and comparative geurotoxicity of the trichothecenes satratoxin G and roridin L2 from Stachybotrys chartarum. J Toxicol Environ Health A, 2009. 72(20): p. 1242-51.
- 28 Jarvis, B.B., et al., Study of toxin production by isolates of Stachybotrys chartarum and Memnoniella echinata isolated during a study of pulmonary hemosiderosis in infants. Appl Environ Microbiol, 1998. 64(10): p. 3620-5.
- 29 Yike, I., T.G. Rand, and D.G. Dearborn, Acute inflammatory responses to Stachybotrys chartarum in the lungs of infant rats: time course and possible mechanisms. Toxicol Sci, 2005. 84(2): p. 408-17.
- 30 Lee, M.G., et al., Effects of satratoxins and other macrocyclic trichothecenes on IL-2 production and viability of EL-4 thymoma cells, I Toxicol Environ Health A, 1999, 57(7); p. 459-74. 31 Thrasher, J.D. and S. Crawley, The biocontaminants and complexity of damp indoor spaces: more than what meets the eyes. Toxicol Ind Health, 2009. 25(9-10): p. 583-615
- 32 Nagase, M., et al., Apoptosis induction by T-2 toxin: activation of caspase-9, caspase-3, and DFF-40/CAD through cytosolic release of cytochrome c in HL-60 cells. Biosci Biotechnol Biochem, 2001. 65(8): p. 1741-7.
- 33 Wu, Q., et al., Trichothecenes: immunomodulatory effects, mechanisms, and anti-cancer potential. Arch Toxicol, 2017. 91(12): p. 3737-3785.
- 34 Schlam, D., et al., Gliotoxin Suppresses Macrophage Immune Function by Subverting Phosphatidylinositol 3,4,5-Trisphosphate Homeostasis. mBio, 2016. 7(2): p. e02242.
- 35 Xiao, W., et al., Sputum signatures for invasive pulmonary aspergillosis in patients with underlying respiratory diseases (SPARED): study protocol for a prospective diagnostic trial. BMC Infect Dis, 2018. 18(1): p. 271.
- 36 Kapoor, T., et al., Forskolin, an Adenylcyclase/cAMP/CREB Signaling Activator Restoring Myelin-Associated Oligodendrocyte Destruction in Experimental Ethidium Bromide Model of Multiple Sciencis. Cells, 2022. 11(18).
- 37 Kowalska, K., D.E. Habrowska-Gorczynska, and A.W. Piastowska-Ciesielska, Zearalenone as an endocrine disruptor in humans. Environ Toxicol Pharmacol, 2016. 48: p. 141-149. 38 Yan, W.K., et al., Zearalenone affects the growth of endometriosis via estrogen signaling and inflammatory pathways. Ecotoxicol Environ Saf, 2022. 241: p. 113826.
- 39 Lo, E.K.K., et al., Low dose of zearalenone elevated colon cancer cell growth through G protein-coupled estrogenic receptor. Sci Rep, 2021. 11(1): p. 7403. 40 Kowalska, K., et al., ERbeta and NFkappaB-Modulators of Zearalenone-Induced Oxidative Stress in Human Prostate Cancer Cells. Toxins (Basel), 2020. 12(3).
- 41 Lee, R., et al., Zearalenone Induces Apoptosis and Autophagy in a Spermatogonia Cell Line. Toxins (Basel), 2022. 14(2).
- 42 Massart, F. and G. Saggese, Oestrogenic mycotoxin exposures and precocious pubertal development. Int J Androl, 2010. 33(2): p. 369-76.



## **Environmental Fungal Count**

Company:Accession #:Collected:09/07/2025Project Name:External ID:Received:09/12/2025Address:Phone:Completed:09/17/2025E-Mail:

**Dust** 

**Procedure Type:** Quantitative PCR (Polymerase Chain Reaction)

Code	Test - Cotton Swab	Value (Fungal Elements / ML)
AALT	Alternaria alternata	0
AFLAV	Aspergillus flavus	0
AFUMI	Aspergillus fumigatus	0
ANIG	Aspergillus niger	5,740
AOCHR	Aspergillus ochraceus	0
APENI	Aspergillus penicillioides	121
ASP25	Aspergillus terreus	0
AVERS	Aspergillus versicolor	0
CAU	Candida auris	Q
CGLOB	Chaetomium globosum	202
FUS14	Fusarium solani	0
PBREV	Penicillium brevicompactum	0
PCHRY	Penicillium chrysogenum	265,219
STAC	Stachybotrys chartarum	0
WSEBI	Wallemia sebi	3,267

RTL maintains liability limited to cost of analysis. Interpretation of the data contained in this report is the responsibility of the client. This report relates only to the samples reported above and may not be reproduced, except in full,

without written approval by RTL. The above test report relates only to the items tested.

RTL bears no responsibility for sample collection activities or analytical method limitations.



MOLD	MYCOTOXIN PRODUCED	POTENTIAL HEALTH ISSUES
Alternaria alternata	Alternariol	Spores can be found in damp, poorly ventilated homes, contributing to sick building syndrome [1]. Alternariol has been reported to be mutagenic [2].
Aspergillus flavus	Gliotoxin, Aflatoxin	A. flavus is the second leading cause of invasive aspergillosis in immunocompromised patients. Particularly common clinical syndromes associated with A. flavus include: chronic granulomatous sinusitis, keratitis, cutaneous aspergillosis, wound infections and osteomyelitis following trauma and inoculation [3, 4]. Can cause liver cancer in humans [5].
Aspergillus fumigatus	Gliotoxin, Aflatoxin	A. fumigatus is frequently found in homes and buildings [1]. It is considered to be an opportunistic pathogen, meaning it rarely infects healthy individuals, but is the leading cause of invasive aspergillosis (IA) in immunocompromised individuals such as cancer, HIV or transplant patients [2].
Aspergillus niger	Ochratoxin, Gliotoxin	A. niger produces gliotoxin, which has been identified in the sera of humans and mice with aspergillosis. Causes immunosuppression in patients [8].
Aspergillus ochraceus	Ochratoxin	Ochratoxin has been demonstrated to be Nephrotoxic, Hepatotoxic, and Carcinogenic and is a potent teratogen and immune-suppressant [8]. It has also been associated with urinary tract infections and bladder cancer [9].
Aspergillus penicillioides	Unknown	A. penicillioides is commonly found indoors [10]. Infection has been confirmed in pulmonary aspergillosis [11] and fungal aneurysm [12].
Aspergillus terreus	Gliotoxin, Citirin	Inhalation of fungal spores, which travel down along the respiratory tract, cause the typical respiratory infection [6].
Aspergillus versicolor	Sterigmatocystin	A. versicolor is one of the most frequently found molds in water-damaged buildings. A. versicolor is known to produce a mycotoxin called sterigmatocystin a potentially carcinogenic and hepatotoxic mycotoxin. It is primarily toxic to the liver and kidneys [7].
Candida auris	Unknown	C. auris can be found in healthcare facilities and can be spread through contact with infected patients and equipment. C.auris can cause blood stream infections, wound infections and ear infections [13].
Chaetomium globosum	Chaetoglobosins	C. globosum is a common indoor fungal contaminant of water damaged homes or buildings. Like Stachybotrys, C. globosum spores are relatively large and due to their mode of release are not as easily airborne as are some other molds [11].
Fusarium species	Fumonisins; Zearalenone	Fusarium can cause superficial infections such as keratitis or onychomycosis in healthy individuals and disseminated infections in immunocompromised patients [12].
Penicillium brevicompactum	Ochratoxin A	Producer of the toxin Ochratoxin <i>A. Fungal</i> particles depend on the relative humidity [14]. Can lead to chronic Rhinosinusitis if breathed in high concentrations [15].
Penicillium chrysogenum	Ochratoxin A	Producer of the toxin Ochratoxin 4. Fungal particles depend on the relative humidity [14]. Can lead to chronic Rhinosinusitis if breathed in high concentrations [16]. High levels are correlated with the development of sick building syndrome [17].
Stachybotrys chartarum	Macrocyclic Trichothecenes	S. chartarum, commonly known as black mold, is highly toxic to humans. Nausea, vomiting, diarrhea, burning erythema, ataxia, chills, fever, hypotension, hair loss and confusion are symptoms in individuals living or working inside Stachybotrys infested homes and buildings [10].
Wallemia sebi	Walleminol	Spores from <i>Wallemia</i> are significant allergens. Inhalation exposure of spores can lead to conditions such as asthma, hypersensitivity, or pulmonary fibrosis [23].

## **REFERENCES:**

- 1 Abel-Fernández, E., et al. (2023). Going over Fungal Allergy: Alternaria alternata and Its Allergens. Journal of Fungi, 9(5), 582.
- Ostry, V. (2008). Alternaria mycotoxins: an overview of chemical characterization, producers, toxicity, analysis and occurrence in foodstuffs. *World Mycotoxin Journal, 1*(2), 175-188.
- Deshazo, R. D. (2009). Syndromes of invasive fungal sinusitis. *Medical Mycology, 47*(s1), S309–S314.
- 4 Hedayati, M. T., et al. (2007). Aspergillus flavus: human pathogen, allergen and mycotoxin producer. Microbiology, 153(6), 1677-1692.
- 5 Nixon, M. W. (1990). Aflatoxin and liver cancer. Lancet, 335(8698).
- 6 Vesper, S., et al. (2007). Quantitative PCR analysis of molds in the dust from homes of asthmatic children in North Carolina. *Journal of Environmental Monitoring*, 9(8), 826.
- Roohani, A. H., et al. (2018), Comparing the Profile of Respiratory Fungal Pathogens amongst Immunocompetent and Immunocompromised Hosts, their Susceptibility Pattern and Correlation of Various Opportunistic Respiratory Fungal Infections and their Progression in Relation to the CD4+T-cell Counts. *Indian Journal of Medical Microbiology*, 36(3), 408–415
- 8 Bui-Klimke, T. R., & Wu, F. (2014). Ochratoxin A and human health risk: A Review of the evidence. Critical Reviews in Food Science and Nutrition, 55(13), 1860-1869.
- 9 Sobel, J. D., & Vazquez, J. A. (1999). Fungal infections of the urinary tract. World Journal of Urology, 17(6), 410-414.
- 10 Visagie, C., et al. (2014). Aspergillus, Penicillium and Talaromyces isolated from house dust samples collected around the world. Studies in Mycology, 78(1), 63-139.
- 11 Santana, M. F., et al. (2020). Confirmed Invasive Pulmonary Aspergillosis and COVID-19: the value of postmortem findings to support antemortem management. Revista Da Sociedade Brasileira De Medicina Tropical, 53.
- 12 Tambuzzi, S., et al. (2019). Fungal aneurism of the right posterior inferior cerebellar artery (PICA). Medical Mycology Case Reports, 26, 25-27.
- 13 Shahardashti, R. V., & Lass-Flörl, C. (2019). Antifungal resistance in Aspergillus terreus: A current scenario. Fungal Genetics and Biology, 131, 103247.
- 14 Reijula, K. (2003). Mycotoxins of Aspergilli exposure and health effects. Frontiers in Bioscience, 8(5), 232-235.
- 15 Bradley, S. F. (2019). What Is Known AboutCandida auris. [AMA, 322(15), 1510.
- 16 Kuhn, D. M., & Ghannoum, M. A. (2003). Indoor Mold, Toxigenic Fungi, and Stachybotrys chartarum: Infectious Disease Perspective. Clinical Microbiology Reviews, 16(1), 144-172.
- 17 Cabrera-Aguas, M., et al. (2022). Infectious keratitis: A review. Clinical & Experimental Ophthalmology, 50(5), 543-562.
- 18 Heutte, N., et al. (2016). Assessment of multi-contaminant exposure in a cancer treatment center: a 2-year monitoring of molds, mycotoxins, endotoxins, and glucans in bioaerosols. nvironmental Monitoring and Assessment, 189(1).
- 19 Murr, A. H., et al. (2012). Some chronic rhinosinusitis patients have elevated populations of fungi in their sinuses. The Laryngoscope, 122(7), 1438-1445.
- 20 Bhavsar, S., et al. (2022). Invasive rhinosinusitis due to Penicillium chrysogenum in an adolescent man with new-onset leukaemia: a diagnostic dilemma. BMJ Case Reports, 15(12), e252265.
- 21 Straus, D. C. (2009). Molds, mycotoxins, and sick building syndrome. Toxicology and Industrial Health, 25(9-10), 617-635.
- 22 Straus, D. C. (2011). The possible role of fungal contamination in sick building syndrome. Frontiers in Bioscience, E3(2), 562-580.
- 23 Zajc, J., & Gunde-Cimerman, N. (2018). The genus Wallemia—From contamination of food to health threat. Microorganisms, 6(2), 46.